
### **Historic, Archive Document**

Do not assume content reflects current scientific knowledge, policies, or practices.



# THE HOME OF PACIFIC STRAINS



HYBRIDIZING GARDENS VETTERLE & REINELT CAPITOLA - CALIFORNIA



Display of Tuberous Begonias in One of Our Greenhouses

### TO OUR PATRONS

VISITORS are cordially invited to inspect our plantings during the season. The Polyanthus Primroses are at their best during March and the first part of April. ... In April and May the first crop of Delphinium seedlings is at its height . . . Breeding stock of tuberous Begonias in our greenhouses is at its best in June and July . . . In August and September over a million seedlings of tuberous Begonias can be seen in full bloom in our lath houses . . . New crops of Delphinium seedlings are brought into flower during August and September . . . We are situated four miles out of Santa Cruz, as can be seen on the map. When coming from Santa Cruz, take the first side road to the right when past city limits and follow it to our nursery . . . If coming from Watsonville, turn left in Soquel, as indicated on map.

### JOIN AMERICAN BEGONIA SOCIETY

Descriptive bulletins on culture of all Begonias are issued monthly. Membership, \$1.00 per year. Write to

### **AMERICAN BEGONIA SOCIETY**

757 ATHENS BOULEVARD · LOS ANGELES, CALIF.

### A WORD OF THANKS


We are indebted to you who have contributed to our success in business and made it possible for us to experiment with breeding and improvement of plants on a larger and larger scale . . . The process of breeding, selecting and eliminating is naturally slow, and no spectacular achievements can be had overnight. The higher the standard, the harder and costlier it is to bring further improvement . . . Our appreciation can best be expressed by giving you each year better value and higher quality for your money than we have been able to do before.

At the Oakland Spring Flower Show we have won the Gold Medal for two consecutive years with our Delphinium Hybrids.

We have also been awarded the Gold Medal for our exhibits of tuberous Begonias at the Golden Gate International Exposition in San Francisco, and the Silver Medal at the World's Fair in New York, besides numerous awards of merit for our Montbretias, Primroses, Ranunculus and Freesias.

### HOW TO GET THERE

REVER



### **TERMS**

Cash with order or approved references. Orders received without cash or established credit will be sent C.O.D.

**SEEDS** . . . Are sent prepaid.

**BULBS**... Please add 25c postage and packing charges on *all* orders under \$2.50.

**PLANTS** . . . *All plants* are shipped express, collect. On orders of less than \$1.50 a packing charge of 25c must be added.

SPECIAL NOTE . . . As the season advances certain types, colors and sizes of tubers are sold out. Kindly state whether substitution or refund desired.

To our California customers: *Please do not forget State Sales Tax.* Not chargeable on out of state orders.

# **BEGONIA TUBERS**

(Delivery January to end of March)

The great majority of our tubers are produced annually from seed and ripen with us approximately the end of December. The size will vary anywhere from less than 1 in. in diameter to as much as 3 in. in diameter. The larger the bulb,

the larger the plant one can expect. However, the size and quality of flowers is not governed by the size of the bulb.

For commercial distribution they are graded into three sizes as shown in the picture below.



#### SEEDLING TUBERS

1. Large Size. 2 in. up in diameter.

This is the most preferred size which produces large plants and if grown in pots requires minimum 8 in. or 9 in. containers.

2. Medium Size. 1½ in. to 2 in. diameter.

Often this size is used for smaller pots, 7 in. size, and will give excellent results for bedding.

3. Small Size. 1 in. to  $1\frac{1}{2}$  in. diameter.

This size will not give large plants the first season but it is satisfactory for bedding purposes or small pot plants. We do not offer these in our retail trade as they require professional experience to bring best results.


### **CUTTING TUBERS**

Nos. 4, 5 and 6
In some hanging varieties and multiflora type we offer tubers produced from cuttings which differ only in form from the tubers produced from seedling plants. Cuttings often have to be grown for two years before they form sufficiently large tubers for sale. The cost of production and additional growing naturally raises the price of propagated varieties.

### SPECIMEN TUBERS

No. 7
Each year we have a limited number of large tubers two to four years old, which have been grown in pots for breeding purposes. These, of course, are the finest Begonias we can offer for show purposes to be grown in containers 12 in. in diameter. We can supply practically all types and colors listed on pages 7, 9 and 11 at \$1.00 each or \$10.00 per dozen.

# **BEGONIA PLANTS**



**SEED**... Seed of all types of tuberous Begonias listed on pages 7 and 9 is available either in straight colors or any desired mixture. To produce high-quality seed only the very finest members in each class are used as parents. After thorough testing they are artificially cross-pollinated, as only by this method are we able to control the hereditary factors and produce seed that is in a class by itself.

Seedling plants of all tuberous Begonias listed on pages 7, 9 and 11 are available during May and June. Of course, no seedlings can be supplied of the named series of basket Begonias or the Multiflora Nana as those are propagated by cuttings only.

These are transplanted twice and hardened off before shipping so that upon arrival they can be planted directly outdoors in permanent location. They will begin blooming by July, forming large flowering specimens during the late summer and early autumn.

Regions with warm summers and long, cool autumns, such as most parts of California, will get better results from seedlings than from tubers. Tubers start blooming earlier and the flowers often burn during the hot summer months. For bedding purposes they are preferable as they will form uniform bed and mass of color continuing deep into the autumn months. They can be shipped successfully to all parts of the Pacific Coast but we cannot be responsible for their safe arrival East of the Rocky Mountains.

### **CULTURE OF TUBEROUS BEGONIAS**

Natural state, Central and South America . . . Planting time, January to May. Flowering time, May to November.

**SEED**... Sow from January to March. Place one or two inches of gravel in flats or seed pans, to insure good drainage, over which place a fairly coarse mixture of two-thirds leaf mold and one-third peat, about one inch deep. Smooth the surface with the same mixture, finely sifted, not more than one-eighth inch deep but do not press it down. It is necessary for the surface to be of a spongy character, so that when the seed germinates, the young roots can get into it. If surface is too fine and packed, the seedlings will often fall over, as the young roots are unable to get in... Place the pans in shallow water until thoroughly soaked up from below, then broadcast the seed. Cover with glass and a sheet of paper or keep it in the dark until germination takes place. Night temperature of 65 to 75 degrees Fahrenheit is necessary for quick germination. Lower temperature than 65 degrees will considerably slow up germination, with poorer results... As

soon as germination takes place, take the paper off and in three or four days lift the glass also, otherwise the little seed-lings will get too spindly. Warm temperature, protection from direct sunlight, and uniform moisture, are absolutely necessary. Even a slight drying out of the surface will be fatal to the delicate young plants. Very gentle overhead watering should be used when necessary. . . . Dampening off will occur only if flats are kept too wet in greenhouse without proper ventilation. Watering with a weak solution of Clorox, about 4 to 6 per cent, we found as best cure and preventive.

**TRANSPLANTING** . . . When the third leaves are developed, transplant one inch apart in flats containing the same mixture as for sowing. No finely-sifted surface is necessary any more. Do not put more than one to one and one-half inches deep of soil in your flats. Deep flats filled with several inches of soil are not necessary. Usually they do not drain well, soil will get sour, and checking of the growth



Upper left: Giant Single. Lower left: Crested Type. Upper right: Narcissiflora Type. Lower right: Frilled Type

### TUBEROUS BEGONIAS

### GIANT SINGLE TYPE

The individual flowers of this group often reach astonishing size of from 8 to 12 inches in diameter; carried on strong, stiff stems. They are very decorative as large specimen plants in tubs or for bedding purposes, where they will form a solid sheet of color.

Colors

White Rose Yellow Scarlet
Pink Salmon Orange Dark Red

### SINGLE FRILLED TYPE (CRISPA)

Perhaps the most adorable of the single types. The large, individual flowers are beautifully frilled and ruffled at the edge.

Colors

Solid Colors—White Yellow Scarlet
Pink Orange Dark Red

Variations-Rose Light Salmon Apricot

Crimson Rose Dark Salmon

Fascination: On cream background, edged in rosy red.

### SINGLE NARCISSIFLORA TYPE

A new form recently developed. The flowers resemble somewhat the trumpet daffodils, hence its name. Daffodil flowered. Very effective as a pot plant.

Colors

Yellow Salmon Shades Orange Salmon Shades
Salmon Shades Red Shades

### SINGLE CRESTED TYPE CRISTATA

The accompanying picture speaks for itself as to the beauty of this type, which we have highly improved in recent years. Dwarf, stocky growth, with masses of flowers, make this type suitable both for bedding and show purposes.

Colors

Pink Shades Crimson Rose Yellow Scarlet Rose Shades Salmon Shades Orange Dark Red

### PRICE OF SINGLE TYPES

**TUBERS**... Delivery, January to March. Large size, 20c each; \$2.00 per dozen; \$14.00 per 100. Medium size, 15c each; \$1.50 per dozen; \$10.00 per 100. **SEEDLING PLANTS** . . . Delivery, May-June. \$1.00 per dozen; \$7.50 per 100. F.O.B. Capitola. **SEED** . . . 50c per packet; 25c per half packet.

### **CULTURE OF TUBEROUS BEGONIAS—Continued**

will be the result. Before they get too crowded, transplant in the same manner again farther apart, until strong enough to be planted out in the open or, if desired, potted up... If planting in open ground, enrich the soil liberally with well-rotted cow or sheep manure and a sprinkling of bone meal. If soil is heavy, add plenty of either leaf mold, peat or sand or a mixture of all, so that the ground will be light and porous. Plant in a shaded position, such as under the trees or north side of the house, where direct sunlight cannot reach them. Keep well watered; fine, overhead sprinkling preferred.

**FEEDING** . . . If a very strong growth is desired, additional feeding can be given when the plants reach approximately four to five inches in height, either by the sprinkling of one-half teaspoonful of ammonium phosphate around each plant or approximately a heaping teaspoonful of cottonseed meal, which should be worked into the soil slightly, not too close to the stem. This dose will usually prove sufficient for the whole season but can be duplicated again later in the summer when the plants are in full bloom, which will prolong the flowering season to some extent.

foliage turns yellow, withdraw the water gradually and when all growth dies down entirely, take out, wash off all soil, taking care not to bruise the tubers, dry in sunlight for a day or two until thoroughly dry, then store in open flats in cool, dry place. See that all particles of the old stem are removed until healthy tissue shows; otherwise, if left on, they will decay and destroy the tuber.

**TUBERS** . . . If early flowering is desired, place tubers during January and February in a warm place in open flats, moisten slightly once in a while and gradually, as they begin

sprouting buds, plant in flats filled with peat, only about three inches apart, according to the size of the tuber, so that they are only slightly covered. Keep uniformly moist but not too wet, in a warm place, well lighted, until three or four inches of growth develops; then plant in pots or permanent position as desired. The front is always where the tips of the leaves are pointing. Often, if dormant tubers are planted in the open ground, especially if it is cold and too wet, numbers of them rot and the planting will be uneven, some coming earlier, some later, and facing haphazard way. Started first in peat, they will develop splendid root systems, which are necessary for developing good specimens.

**POT CULTURE** . . . For pot culture, any rich, light soil is suitable. Mixture of two parts of coarse leafmold, one part of loam will bring good results. It is important for mixture to be of light, coarse character, so that it will drain well. Finelysifted soil will pack down without permitting the circulation of air, hence only a moderately good result can be expected. Perfect drainage is most important, and to provide it, place at least one inch deep of gravel on the bottom of the pot. When potting up, use smaller pots in beginning and move later into larger pots before they are pot-bound. If tubers are well started in peat, with considerable root system, they may be planted immediately in large pots approximately from six to eight inches for one-year-old tubers. Older, larger tubers will, of course, require larger pots. Very great care with watering is necessary when they are planted immediately in large pots. Keep the surface only moist in the young stage, until the growth is well developed, and then heavier watering is justified. If you over-water the soil in the beginning, it will become sour, the plants will make only a sickly growth, buds will fall off before blooming and the plants may die out entirely. This is true especially of hanging basket types. If very



# TUBEROUS BEGONIAS

**DOUBLE CAMELLIA TYPE** . . . The largest and most popular of the double types. Individual flowers, from 5 to 8 inches in diameter, resemble camellias and roses in all forms and variations of color.

#### Colors

Solid, Uniform Colors, with slight variation only-

White Crimson Rose Cardinal Red Yellow
Pink American Beauty Scarlet Orange
Rose Salmon Red Dark Red Flame Orange

Apricot

Salmon Shades, varying more or less within themselves—

Blush and Pink Salmons Apricot and Orange Salmons

Yellow and Apricot Salmons Salmon Rose

was developed by crossing the large Camellia with the small Marmorata type. The flowers are slightly smaller but of beautiful Camellia form, with two-tone combinations of color strongly pronounced in a definite contrasting edge.

#### Colors

Red Picotee Rose Picotee Apricot Picotee Salmon Picotee

**FIMBRIATA PLENA TYPE** ... Double frilled, or often called Carnation type. This group has been enormously improved in the last few years, the flowers reaching in some cases the size of the Camellia type with more and more refined form. Its strong, bushy growth makes it exceptionally desirable for bedding and pot plants.

#### Colors

Solid Uniform—White Yellow Orange
Scarlet Dark Red Crimson Rose Salmon Red
With Variations—Blush Pink Rose
Apricot Salmon

**DOUBLE ROSEBUD TYPE** ... Flowers of the most exquisite form, resembling perfect rosebuds, with beautiful, two-tone, pastel-shade coloring. Very tall, sturdy growth makes it especially desirable for growing large specimen plants in tubs for decorative purposes.

#### Colors

Pastel Shades—Blush Pink Rose
Flame Salmon Light Salmon Dark Salmon

### PRICE OF DOUBLE TYPES

TUBERS . . . Delivery, January-March.

Large size: 25c each; \$2.50 per dozen; \$18.00 per 100. Medium size: 20c each; \$2.00 per dozen; \$14.00 per 100.

SEEDLING PLANTS . . . Delivery, May-June. \$1.50 per dozen; \$10.00 per 100. F.O.B. Capitola.

SEED . . . \$1.00 per packet; 50c per half packet.

### **CULTURE OF TUBEROUS BEGONIAS—Continued**

strong growth is desired, it will be necessary when the plants are established in the final pots and have at least four of the leaves developed, to water once a week with liquid manure.

FEEDING . . . A number of quick-acting, commercial fertilizers can also be used. One of the safest is cottonseed meal, which we are using now almost exclusively, as it gives very good results. Half a teaspoonful, once or twice during the season, sprinkled around the edge of the pot (not too close to the stem) will be sufficient. Heavier doses will do more harm than good, often causing distortion in the flowers or burning up the roots, destroying the plant entirely. Lath houses with strong muslin roofs are a very suitable place for development of show specimens in warm regions where greenhouses would be too warm. If grown in a greenhouse, thorough ventilation day and night must be available. Openings in the walls, under the benches, will give good circulation of air. On hot days, heavy shading and sprinkling the walks with water will keep the temperature down. 65 to 80 degrees Fahrenheit is the most ideal temperature, for as the cooler and slower they develop, the finer and larger the flowers will be. Higher temperatures produce quick, weak growth, and the flowers do not develop as good size and as lasting quality.

**PRUNING**... Do not prune Begonias. Any open wound will be immediately attacked by fungus, especially when

crowded, without good air circulation. If you cut the flowers, cut only half of the stem. The other half will mature and fall off itself without leaving a wound.

**PESTS**... Occasionally Begonias are attacked by aphis and thrip in the greenhouse, if kept in dry, close temperatures. Fumigating with nicotine on two or three alternate evenings will control these pests easily. Very rarely they are also subject to mite attacks. These are very small and cannot be seen without a lens. The first signs of their presence are brown, rusty streaks on the young shoots and foliage, deforming and destroying the season's growth entirely if not checked. Infected plants should be isolated immediately. Thorough spraying with Volck will check this pest.

**FAILURES** . . . The most common failure in Begonias is the falling off of buds before they develop, which is caused either by the plants being grown in too small pots without a sufficient amount of food, severe drying out or severe over-watering. In hot weather the flowers develop too rapidly and the plants, by drawing much more water from the soil, tend to throw off the buds. Abundant growth, with little or no flowers, is the result of too heavy shade. Curled and shiny foliage is a sign of too strong an exposure to the sun. The most damage is done by over-feeding; the first signs of it are a soft, glassy texture of the foliage, curling under gradually, wilting and dying off.



# **TUBEROUS BEGONIAS**

### DOUBLE HANGING TYPE

Perhaps the most attractive type of tuberous Begonia. The hanging habit, with great masses of blooms, makes them favorite for decoration of greenhouses, open verandas or sun porches where they can be protected against a strong light and wind.

**SEEDLING TUBERS** . . . We have made a large amount of experimental crosses and are offering great variety of new color combinations and forms which were grouped in the following shades.

Varieties with large flowers . . .

White Blush Shades Pink Shades

Yellow Shades Orange Red Shades

These are second generation crosses between the double Camellia and Hanging type and are suitable chiefly for displaying large specimen plants. They do not have as ideal hanging habit as the smaller types. However, if pinched so they can branch they will form beautiful hanging specimens.

Varieties with medium sized flowers . . .

These have greater profusion of bloom and better hanging habit.

White Blush Shades Pink Shades

Rose Shades Crimson Rose Shades Salmon Rose Shades

Scarlet Shades

### PRICE

**TUBERS**... Delivery, January to end of March.

Large size, 25c each; \$2.50 per dozen; \$18.00 per 100.

Medium size, 20c each; \$2.00 per dozen; \$14.00 per 100.

SEEDLING PLANTS... \$1.50 per dozen; \$10.00 per 100.

F.O.B. Capitola

### NAMED VARIETIES

We have been selecting and propagating a number of seedlings of the basket Begonias which are outstanding in color, form of flowers and hanging habit. All are propagated by cuttings, forming rather irregular tubers in comparison to the seedling tubers which are smooth and round. . . . It takes several years to propagate a sufficient amount of tubers for introduction and we are so far able to offer only the two following varieties.

*Pink Shower* . . . The individual flowers are a delicate cerise-pink, holding the beautiful form to the very end of the season in such profusion that the plant portrays the name it bears exactly.

Avalanche... Recent introduction and a good companion to Pink Shower, as it is almost a white counterpart of it. A slight greenish tinge in the center of the flowers gives it added attraction.

PRICE . . . 75c each; \$7.50 per dozen.

**SEED** . . . To produce a high quality seed of the Hanging type is very difficult and we are often not able to supply all colors that we list. Each year we make entirely new crosses to improve either form or hanging habit or derive new colors. This year we can offer only a small quantity in the following shades.

White Pink Rose Salmon Shades Red Shades

Complete mixture of the above and all new crosses made this season.

**PRICE** . . . 50c per half packet; \$1.00 per packet.

### **MULTIFLORA NANA**

This group of tuberous Begonias which we are offering this year are not exactly new. However, they have not been distributed or grown in America with the popularity which they rightly deserve, as they have been in Europe for the last twenty-five years. They differ from the large-flowered types in having a dwarf, bushy growth covered all summer with a mass of small double flowers 1½ to 3 inches in diameter. The character of their growth makes them invaluable for borders or bedding purposes, as they form a veritable carpet of single color. All varieties are propagated artificially, by cuttings only, as seed from them does not come true to type or color. In this manner the bulbs offered are divisions of the original plant and members of individual variety, forming absolutely no variation.

Fireflame . . . Flamboyant. Small semi-double flowers of the most intensive scarlet. The plants grow only four to six inches high and are simply covered with blooms from June to the end of October. It can be utilized both for bedding and hanging baskets.

**PRICE** . . . 25c each; \$2.50 per dozen; \$18.00 per 100.

Mrs. Helen Harms . . . Pure yellow double. A very dwarf, bushy growth; outstanding for borders or bedding.

**PRICE** . . . 35c each; \$3.50 per dozen; \$25.00 per 100.

Mme. Richard Galle . . . Pure orange double. Almost identical with Mrs. Helen Harms, except for color, which is clear orange.

**PRICE**... 35c each; \$3.50 per dozen; \$25.00 per 100.



### SPECIAL . . . CHOICE SELECT VARIETIES OF TUBEROUS BEGONIAS

During flowering season we select and mark all the choicest new seedlings which are above average in size, form and color. We are offering tubers of these to connoisseurs who desire only the finest for their gardens. For show purposes and greenhouse decoration they are the last word in tuberous Begonias. In this group we supply all the double types and colors listed

on the previous pages, besides including also all the new color developments derived from our crosses preliminary to introduction. The double Camellia "Apple Blossom" shades are supplied in this group only.

**PRICE**... Tubers, delivered January to March. Large size, 50c each; \$5 per dozen; \$37.50 per 100. Medium size, 40c each; \$4 per dozen; \$30 per 100.

# BEGONIA HIEMALIS

(Winter Flowering Socotrana Hybrids)

An entirely new group of tuberous Begonias developed in England by crossing B. Socotrana with largeflowered tuberous Begonias. Their cultural requirements are slightly different, yet simple. Their chief asset is the flowering season, which can be extended from October to the end of January. The fact that they will grow very well in the house will insure them extreme popularity. The accompanying picture was taken after the plant was displayed three weeks on a mantle shelf in a living room in December. All varieties in this group have to be propagated by cuttings, as the plants do not form seed. The cuttings form tubers in their second year. We are in a position to offer a limited number of tubers during March and April, when they have their resting season.



#### NAMED VARIETIES

Conquest . . . A very strong grower with dark green foliage, bearing large trusses of single orange flowers shaded slightly with rose.

Clibrans Red . . . Perhaps the best of the single red varieties, the color being ox-blood red.

*Emily Clibran* . . . Large trusses of medium-sized, double flowers, edged deeper apricot, on a creamy apricot ground. Very rapid grower and good increaser.

Lady Roberts . . . Creamy white, double flowers, showing green tinge in the center and turning blush-salmon with age. One of the strongest growers and most rapid increaser.

**PRICE**... Above four varieties, 50c each; \$5.00 per dozen. *Fascination*... The most outstanding of all single varieties, simply covered with clear, apricot-salmon flowers. Dwarf, bushy growth; makes excellent pot plants.

Rose Queen... The form of the flowers resembles that of the Sweet Pea and the whole plant looks like a single bouquet when in bloom. The color is clear, deep rose.

**PRICE** . . . \$1.00 each; \$10.00 per dozen.

### UNNAMED SEEDLINGS

We have been experimenting with crossing the Begonia Socotrana with the best of our tuberous Begonias and after several years of trying have been successful in securing a small quantity of seed. The plants from this seed are in full bloom this autumn and we were rather nicely surprised that the average quality was far above most of the named varieties that we imported previously from abroad. Only a few of the very most outstanding will be propagated, named, and eventually introduced when available in quantity. In the meantime we are offering large tubers of the mixed seedlings which will perhaps give much finer results than anything we offered previously. All of them are double, with medium large flowers and very strong growth.

*Orange-Scarlet Shades* (Camellia Flame Orange x Socotrana) . . . All of these are of uniform color and type of growth, varying only slightly in form and size of flowers.

Mixed Shades (Various Tuberous Begonias x Socotrana) . . . The color variation in this group is limited to Rose, Crimson-Rose, Salmon-Rose and Dark Red. The form varies according to the parent used and as these crosses were made in small quantity only, they were combined and are sold only in a mixture. Delivery, March-April.

**STRONG TUBERS** . . . 50c each; \$5.00 per dozen; \$35.00 per hundred.

[ PAGE THIRTEEN ]

### BEGONIA HIEMALIS

(Winter Flowering Socatrana Hybrids)



### **CULTURAL DIRECTIONS**

**SOCATRANA HYBRIDS** . . . The culture of the Socatrana hybrids differs slightly from the summer-flowering tuberous Begonias. Originally they were produced by crossing Begonia Socatrana with tuberous Begonias. All the offsprings are sterile and have to be propagated artificially by cuttings.

PROPAGATION BY CUTTINGS . . . Young shoots, formed in the axils of the leaves in April and May when the resting period is over, may be cut when approximately two to three inches long with the basal ring and rooted in sand. If bottom heat is available, they will root within two or three weeks. Under cooler conditions they will root a little slower but very freely. They should be kept close and heavily shaded but as soon as they begin rooting, ventilation should take place. Potted up in three-inch pots, they will begin growing freely if kept either in the greenhouse or out-of-doors in a lath house. Before becoming pot-bound they should be transplanted into larger pots, as necessary during the summer, and pinched once or even twice to prevent them from blooming too early and to form a nice, bushy growth. In this manner cuttings rooted in June will require approximately six- to seveninch pots by September. Those rooted in July will finish in smaller pots.

FLOWERING SEASON . . . By October the plants will commence to bloom and will continue to do so throughout the winter, if kept at a temperature of 60 to 65 degrees, which suits them best during the flowering season. When the flowering period is over, the plants rest from the end of January until the end of March. At this time water should be given very carefully, just sufficient to retain the stems and leaves; if too much is allowed, rot may set in. On the other hand, if insufficient water is given, the stems and leaves fall away and the plant is much less ready to commence growth when required. Early in April, when growth recommences, water can be given more

HOUSE CULTURE . . . Those who have no greenhouse facilities may grow these Begonias in pots out doors well protected against the sun but giving sufficient light to prevent them from being spindly. When cool weather approaches in October they may be transferred into the house where if given sufficient light they will bloom very freely, especially if kept in a cool, dry temperature.

**TUBERS** . . . When plants attain an age of two years the tubers are sufficiently large to shake out of the soil for a resting period. These can be restarted in May in a mixture of coarse sand and leafmold and potted up later as described for cuttings. Several shoots will appear on the tuber, some of which may be cut off with the basal ring and rooted in the same manner as the shoots from leaf axils. Individual leaves can be cut off and rooted in sand but the percentage of them forming plants is comparatively small and varies with the variety.

**SOIL**... The mixture best suited for these Begonias consists of two parts of loam, two parts of leafmold and one part of sand. A sprinkling of bone meal added will be beneficial. For the final planting one part of well rotted manure can be incorporated into the soil. Always see that the soil mixture is coarse and light, and pot lightly at all times.

**PESTS**... Rusty foliage is a sign that the mite is present, or a species of thrip. Both can be exterminated by fumigation or spraying with nicotine. Aphis will attack some of the varieties under greenhouse conditions; these too can be exterminated in the same manner. Syringing twice a day during the summer growth will keep these pests down if grown out of doors. Under greenhouse conditions keep the temperature quite moist with good ventilation, seeing that the foliage is always dry before evening, otherwise rot may take place.

### EARLHAM STRAIN OF GIANT MONTBRETIAS

We have imported, directly from the originator in England, a collection of some forty varieties of the new hybrid Montbretias, which under good conditions reach the height of four feet, with individual flowers from two to three inches in diameter. They require slightly richer soil and better cultural conditions than the old type of Montbretias but will repay in full measure for the extra effort. Their brilliant red-gold and copper tones make a splash of color for six weeks during the summer. Excellent as cut flowers, lasting fully two weeks in water. The five following varieties, which we are introducing for the first time in the United States, are the most outstanding and distinct of the entire collection. They are fully as large as His Majesty.

	Price	23	
Cecil: Buff-yellow, with apricot eye; large flowers; very tall, four feet or over	Each .20	Dozen \$2.00	Hundred \$15.00
Jessie: Equally as fine as Cecil but a shade deeper in coloring		2.00	15.00
Lord Lambourne: An improvement over Pocohontas, which we used to list, the color being a deeper shade of bronzy-orange-scarlet, with more graceful sprays and taller stems	.20	2.00	15.00
Sir A. M. Wilson: A uniform shade of crimson, the deepest color so far achieved in Montbretias. It has a splendid type of growth, with graceful	.20	2.00	15.00
J. A. Fitt: A masterpiece of color, which is a brilliant orange-scarlet self. It is easily the finest Montbretia in existence, named after its originator. It received a first-class certificate from the Royal Horticultural Society in England	.30	3.00	20.00
His Majesty: The well-known and still one of the most outstanding Montbretias, which is available now at a reasonable price. The flowers, which reach three inches in diameter, are carried in graceful sprays on four-foot stems. Their copperyorange color blends well with most summer			
flowers	.10	1.00	7.00

CULTURE OF MONTBRETIAS . . . The new strain of Earlham Montbretias requires somewhat richer cultivation than the old, small-flowered types of Montbretias. The bulbs are distributed from December to March and should not be kept in a dry state longer than necessary. As a rule, under California conditions, if transplanting is desired they can be dug up in autumn and replanted immediately. In a region where deep frosts prevail, it is advisable to lift them after the frost has cut down the foliage, leaving the soil adhering to the roots, and store them in the basement, where they can be kept thoroughly dry until spring, when planting is again possible. For good results, the soil should be dug up quite deeply, well enriched with bone meal and old, well-rotted manure, before planting. Fresh manures, if used, are often more harmful than beneficial. No additional feeding is required during the growth but a continuous supply of moisture is necessary until the flowering season is over, after which time they can be kept on the dry side so the tubers will be well ripened when the time for lifting comes. They multiply very rapidly if grown in light soil. In heavy soils they will produce strong shoots but the increase is much slower.



Galahad Series

### PACIFIC STRAIN OF

Although a comparatively short time has passed since our strain of Delphinium was introduced to the gardening world, the enthusiastic acclaim of amateurs and professionals alike has made us feel that we have contributed a step in the progress of better Delphiniums. At 1939 Oakland Spring Garden Show, which is the annual flower event of the Pacific Coast, our Delphiniums received the gold medal for the most outstanding display in the show, in open competition with all other spring flowers. These plants were field-grown seedlings in their maiden bloom, dug and balled the day before the show, without being forced in any way. We were again awarded the gold medal in 1940, the picture of our exhibit being reproduced on the back cover of this catalogue.

Contrary to the popular practice of naming individual plants, our aim is to produce a group of individual plants resembling each other as closely as possible, keep interbreeding them until the progeny brings a high percentage of specimens true to description and then introduce it as a named series. In this manner we can offer new progressions of each series each year as the progeny always supersedes the parents. The names of our Delphinium series, with the exception of the blues, which have descriptive names, have been chosen from Tennyson's "Idylls of the King."

### **NEW INTRODUCTION**

Blue Bird Series ... A true blue Delphinium has been the hardest thing to produce and we have been reserving the name Blue Bird to signify our best achievement in this color. We can safely recommend this series as being the clearest medium blues with white bees. They not only come true to color but carry well formed, round flowers  $2\frac{1}{2}$  inches in diameter, on very long, graceful spikes. The habit is all one could desire in modern Delphinium and this series sets a new standard in blues of today.

**PRICE** . . . Cross-Pollinated Seed: \$5.00 per packet; \$2.50 per half packet.

#### RECENT INTRODUCTIONS

Cameliard Series . . . A clear lavender self with white bee, possessing classical perfection of form of the individual flowers, which are very round and attain very large size, averaging 3 inches in diameter. Beautifully balanced spikes, combined with excellent growing habits.

Galahad Series... Our greatest achievement in giant whites, fully worthy of its name, which possesses all the qualities one expects to find in a fine Delphinium. The standard attained in this series is perhaps responsible for the wide popularity it met, ranking in demand next to the blues. The clear white flowers, with white bees, are of large size, reaching often 3 inches in diameter, combined with glistening, heavy texture.

Round Table Series . . . As the name implies, this represents all of the color combinations of the Knights and their Ladies which will eventually be introduced. This year some three hundred different crosses were used to make up this group, combining all the newest color developments and possessing size that will make them invaluable for show purposes.

**PRICE**... Cross-Pollinated Seed: \$3.00 per packet; \$1.50 per half packet. Collection of three packets, \$8.00; collection of three half packets, \$4.00.

[PAGE SIXTEEN]

### **DELPHINIUM HYBRIDS**

Black Knight Series . . . The darkest violet. Individual flowers 2½ to 3 inches in diameter; of beautiful round form, with heavy velvety texture, which gives the color luminosity and vividness not seen before in Delphinium. Very long, well-formed spikes; a true show flower with black bees.

Blue Jay Series . . . Every Delphinium lover seeks the true blue color, which is so rare in nature and so difficult to obtain. These series are our bid toward attaining it. The color is striking, clear medium to dark blue; very intense and alive, with dark, contrasting bee.

King Arthur Series . . . This is still one of the most brilliant Delphiniums we have so far developed. The color is a rich, royal purple, with a velvety texture and large white bee, and the beautifully-formed, long spikes are carried on thin, woody stems.

Guinevere Series . . . A clear, pink-lavender self, with white bee; with very large individual flowers averaging up to 3 inches in diameter. This is a true exhibition type and a beautiful garden plant. Summer Skies Series . . . As the name indicates, this is the blue of a summer sky, with white bees representing fleecy clouds; a color long sought in the hybrid Delphiniums.

**PRICE**...Cross-pollinated seed . . . \$2.50 per packet; \$1.75 per half packet. Collection of five packets, \$10.00; collection of five half packets, \$5.00.

### STANDARD COLORS

Besides the named series we are offering several groups which are not yet segregated to such a degree that they would come true to color. They are the best we have been able so far to produce in their respective colors and are fully equal in character of growth and size of flowers to the named series. Produced by artificial cross-pollination to intensify the desirable characteristics in the highest degree. It is the same seed we use for our own planting, and by releasing it to the public we feel that we are giving them an equal chance. The percentage of show specimens from this seed is the highest obtainable.

Lavender Shades . . . Clear lavender mauves, with white bee.

Lavender Pastels . . . Lavenders with smoky tinge in the petals and dark brown or black bee.

Light Violet Shades . . . Great variation of light violet tones, ranging from lilac to deep violet.

Dark Violet Shades . . . Comprising the deepest color tones in Delphinium, mostly dark bees.

Light Blues . . . Ranging from light to medium blue with white, fawn and dark bees.

Medium Blues . . . Clearest blues are in this group, with white and dark bees.

Dark Blues . . . Ranging from medium to very dark blues, predominantly white bees.

Blue Violet Shades ... Richest coloring in Delphinium comprises this group. Combination of dark blue and violet is further intensified with dark bees.

Clear White . . . These are slightly different from the Galahad Series, the flowers not being as large; however, they are the purest white, with velvety texture. Very heavy branching and long trusses of flowers make them ideal for cut flower purposes.

White—Black Bee . . . The best whites with dark bees we have been able so far to produce.

**PRICE**.. \$2.00 per packet; \$1.00 per half packet. Collection of five colors, \$8.00; half packets, \$4.00. Collection of ten colors, \$16.00; half packets, \$8.00.

[PAGE SEVENTEEN]



Blue Jay Series



King Arthur Series

### SELF-POLLINATED SEED

The qualities we look for in Delphiniums are not only huge flowers and spikes regardless of other characteristics, but on the contrary we often use plants of smaller size if they have outstanding qualities in color or form. Our aim is to produce large flowers properly proportioned to the size of spikes, so as to maintain balance and retain the grace of the entire plant. Another important feature in our developments is comparative mildew resistance. Over ten acres of seedlings are being grown this year for selection, from which only the finest members combining all the desirable characteristics are retained for seed production. This enables us to produce very high quality seed, giving customers the advantages of the newest developments.

The self-pollinized seed offered on this page is collected only from the outstanding plants and it will bring high uniformity of type and color as described below.

#### NAMED SERIES

(For description see page 16)

Blue Bird Series . . .

**PRICE** ... Per packet, \$2.00; per half packet, \$1.00.

Cameliard Series . . . Galahad Series . . . Round Table Series . . .

**PRICE** ... Per packet, \$1.50; per half packet, 75c.

Black Knight Series . . . Blue Jay Series . . . Guinevere Series . . . King Arthur Series . . . Summer Skies Series . . .

**PRICE** . . . Per packet, \$1.25; per half packet, 65c.

#### STANDARD COLORS

Lavender Shades ... Lavender Pastels ... Light Violet Shades ... Dark Violet Shades ... Light Blue Shades ... Medium Blue Shades ... Dark Blue Shades ... Blue Violet Shades ... Clear White and White-Black Bee.

**PRICE** ... Per packet, \$1.00; per half packet, 50c. Collection of five: \$4.00; half packets, \$2.00. Collection of ten: \$7.50; half packets, \$3.75.

### **DELPHINIUM PLANTS**

From March to the end of June and from September to the end of November we are offering strong, young plants in twelve assorted shades as marked for cross-pollinized seed. Transplanted, they will grow rapidly on and bring far better results in a shorter time than the old clumps. We prefer distribution of young seedlings rather than the one- or two-year-old clumps, which very seldom bring good results, as they are quite exhausted and it takes a year at best to get them re-established, during which time the flowers are not at their best. Young seedlings will produce show specimens within four or five months after planting and continue blooming for several years. In California interior valleys where the climate is too warm, autumn planting, September and October, is preferable. Plants having time to establish good root systems during winter will bring beautiful spikes in early spring. Cooler regions will give good results from both autumn and spring planting, except in heavy, cold soils, where spring planting is preferable.

**SEEDLING PLANTS**... \$2.00 per dozen; \$15.00 per 100. F.O.B. Capitola.

[PAGE EIGHTEEN]

# CULTURE OF DELPHINIUM HYBRIDS

SEED . . . Delphiniums can easily be grown from seed which can be sown practically any time of the year, according to the climatic conditions, equipment available or time of flowering desired. Under California conditions, for early spring flowers, sow from June to September; midsummer blooms, December to January; and for fall blooms, February to April. In regions with severe winters, the early summer sowing, from June to July, is recommended, so that the little plants are well established before the winter sets in. Otherwise, the most practical time to sow is early spring. Seed can be kept in good condition for several years if kept in air-tight containers in a refrigerator. In fact, it is practicable even with fresh seed to place it between two moist blotting papers and leave it directly under the freezing compartment of the refrigerator for at least a week to induce higher germination. For best results, sow in flats in a mixture of two-thirds coarse leaf-mold and one-third loam, covering slightly with the same mixture, and moisten thoroughly. To prevent evaporation, cover the flats with newspaper and glass until germination takes place. Immediately after the young plants begin to appear, both the glass and the newspaper should be taken off; however, the plants should be kept shaded and kept moist constantly.

**GERMINATION** . . . If the seeds do not germinate 100 per cent and the flats are exposed to strong light immediately after the first few plants have appeared, the rest may not germinate at all. Artificial heat can be used for germination in early spring and it will give far better results than the natural heat in summer. To get the best results in germination, bottom heat is necessary, with a cool temperature overhead. This is well supplied under glass in early spring; however, in summer, due to hot weather conditions, it is usually the opposite and this is why many people have failures even with the very freshest seed. At temperatures of 85 degrees Fahrenheit and up, the germination is often very poor and what germinates stands a very good chance of being simply cooked. To prevent this, after the seeds are sown in flats, place them on the floor in a cool room or shed, where they can be kept dark for the first ten days, until germination takes place; then give light and fresh air, but keep them well protected against any drying out. We have repeatedly checked on seeds which were sown under greenhouse conditions in summer or the cool shed method or a well-shaded lath house, and the results in each case were from 20 to 50 per cent better under lath house conditions than under the greenhouse conditions. Under slow, cool germination practically every seed will gradually come up. Under quick, hot conditions only a few will germinate, the rest remaining dormant in the soil. Dampening off will occur only if flats are kept too wet in the greenhouse, without proper ventilation. Watering with a weak solution of Clorox, 2 to 4%, we have found to be the best cure and preventive.

**TRANSPLANTING** ... When second leaves are developed and before plants are too crowded, prick in flats three inches apart in a mixture of two-thirds sandy loam and one-third leafmold. Keep shaded for two or three weeks; then gradually give more light and, when larger, harden off in full sunlight before planting out in permanent position about two or three feet apart. Open, sunny location is necessary for best development. In shade they will grow too spindly, with only small flower spikes. If too close to walls, they will mildew more than in the open. The ground should be well prepared for planting. A liberal application of well-rotted cow or sheep manure, with a sprinkling of bone meal mixed with the soil, which should be dug a foot deep, will produce fine growth. Heavy, wet soils will require a slight addition of lime. Good drainage and uniform supply of moisture during the growth is essential.

**FEEDING** . . . When the first crop of blooms has faded, cut the flower spikes off just above the foliage and keep slightly dry for two or three weeks, to give the plants time to rest before the new shoots appear above the ground. When this takes place, cut the rest of the old stock off, sprinkle a teaspoonful of ammonium phosphate around each plant, rake it into the soil slightly and water thoroughly. From the new shoots appearing from the ground select two or three of the strongest and break the rest out. The remaining ones will develop into fine spikes again. Do not force a new growth late in autumn; rather keep the plants on the dry side, because if forced into bringing a third crop late in the season the plants will soon exhaust themselves and gradually die out.

**DISEASES** . . . Mildew will attack plants grown close to a wall, or planted thickly together, especially later in the fall. The new "Ortho Mildew Spray," originated by the California Spray-Chemical Corporation, Richmond, California, diluted with water to a 4 per cent strength, will control it perfectly. This spray not only acts as a preventive but will cure heavily-affected plants, as it kills the mildew instantly. It is a light oil and can be applied at any time without burning the foliage, to which it gives a nice, shiny appearance.

# JOIN AMERICAN DELPHINIUM SOCIETY

Beautiful Yearbook containing history of the newest developments in the Delphinium world. Published annually. Membership, \$2.00 per year.

 $Write\ to$ 

THE AMERICAN DELPHINIUM SOCIETY Ridgefield, Connecticut

Pacific Strain of Polyanthus Primroses

### PACIFIC STRAIN OF POLYANTHUS PRIMROSES

As the Polyanthus group is the most hardy and easiest to grow of all the Primrose family, succeeding in practically every climate, we have concentrated our work on improving this particular type. Our aim is to develop a wider and clearer color range, taller stems, and larger size of individual flowers and flower heads. After testing all of the important strains in commerce, we spent years of breeding and selecting from some 20,000 seedlings each year. We introduced our strain experimentally at the Golden Gate International Exposition on Treasure Island, where the exhibit, although comparatively small, created such interest that our stock was completely sold out within two weeks. This is, perhaps, the best recommendation our strain could have.

**SELECTED PLANTS** . . . We can supply, during August and September, strong divisions of our breeding stock from the previous season, which are the most outstanding developments we have so far produced.

Colors

White Scarlet Golden Yellow
Carmine-Rose Dark Red Bronze-Apricot

Orange Pastels New Blends

**PRICE** . . . 25c each; \$2.50 per dozen; \$15.00 per 100. F.O.B. Capitola

**SEEDLING PLANTS** . . . During February and March we ship strong seedling plants with flowers in assorted colors as per selected plants.

**PRICE** . . . \$1.50 per dozen; \$9.00 per 100. F.O.B. Capitola

cross-pollination from the most outstanding new seedlings each year. By this method we can offer a quality which is in a class by itself. The production, naturally, is very limited and very costly but only in this way can the finest results be obtained. The new crop is available by July 1, which is also the best time for sowing. We pack approximately 125 seeds to a packet which, under reasonably good conditions, will germinate within two weeks after sowing. All colors listed as per selected plants are available separately or in complete mixture.

PRICE . . \$1.00 per packet. (All seed sold out until July, 1941.)

### RICHARDIA ELLIOTTIANA

(Yellow Calla Lily)

Beautiful summer flowering plant of simple cultural requirements. Planting time February to end of April. Strong bulbs from November to February.

**PRICE** . . . 25c each; \$2.50 per dozen; \$18.00 per 100.



#### CALLA AETHIOPICA

(White Calla Lily)

Planting time July-August . . . The white Calla Lilies require swampy conditions for the best development. The flowering season is during winter months and early spring.

**PRICE**... Strong tubers 25c each; \$2.50 per dozen; \$18.00 per 100.

[PAGE TWENTY-ONE]



# RANUNCULUS

We have been inter-breeding the Tecolote and the Australian strain which are the foremost of today, and are offering select stock resulting from these crosses. All single plants are weeded out, so that only uniform, selected plants are offered in the following color groups.

**TUBERS**... White and Cream Shades, Yellow and Gold Shades, Apricot and Orange Shades, Scarlet and Dark Red Shades, Blush, Pink and Rose Shades. *Complete Mixture*. Delivery, July to November.

**PRICE** . . . Large Tubers, \$1.00 per dozen; \$7.00 per 100. Medium size, 75c per dozen; \$5.00 per 100.

**SEED** . . . Ranunculus can readily be grown from seed, especially under California conditions. If sown during September or October and transplanted in December to permanent place, they will produce flowering specimens during March and April. In colder regions, only spring sowing can be recommended. Seed gathered only from choicest specimens can be supplied in separate colors as described, per tubers, or in complete mixture.

PRICE . . . 50c per packet; \$15.00 per ounce.

[PAGE TWENTY-TWO]

	erent from ma		ORDER	BLA	NK	Please do i	
dress)		JATOT MEMAN	o: VETTER	LE & REII	NELT BEE	No	WHITE AND A STATE OF THE PARTY
			HYBRIDIZI	Rec'd			
EASE WRITE OR PI	RINT PLAIN	NLY	CAPITOLA,	Filled			
nd to							
dress						When	n to Ship
		KINDLY LIS	ST TUBERS, PLAI	NTS AND SEE	EDS SEPARATELY		
QUANTITY KIND OF TUBERS		SIZE COLOR PRICE EACH			TOTAL PRICE	SECOND COLOR CHOICE	
							20
ADD ST ERROR		PATOT	35181	4 2100		TO CHO	WHISTANA
23,045		41.469	The same of the sa				
				1921	-	200	-
	4						
	- W	rolling.					-
		1					
						The second second	4 4 1 1 1
The same of the sa			- The state of the				
			3000068	NO AND DE			
ow remittance made up:	September		Principle Plants	A CANADA II			
eck			California Stat		\$		
sh			Packing, etc.				
The second secon			Total amount	, nes andres	sioli-iss	and adalieve on	
oney OrderPlease do not remit by sta	amps				The second secon		

SPECIAL NOTE—As the season advances certain types, colors and sizes of tubers are sold out. Kindly state whether substitution or refund desired.

#### PLANTS

QUANTITY WANTED	KIND OF PLANTS	SIZE	COLOR	PRICE	TOTAL PRICE	SECOND COLOR CHOICE
W. T.	- Jan 1990		ZMBOMAS ABROGAL	MISTOLATION S		
	the seatth of					ALLESS MOJE OF EUDALLY
						The second secon
0.42-4	ned V7					ecololis
	1	112113813	20232 008	ETMAJŠ . SW	TOTAL PROPERTY.	
*15	GO GMODE!	13 (30) (30) (30) (30)	12 S			RC SPRING WYTELYAUG

### SEEDS

QUANTITY WANTED	KIND OF SEEDS	SIZE	COLOR	PRICE EACH	TOTAL PRICE	SECOND COLOR CHOICE
			1,34,54			
			Was a second			
		D-12-4				

#### PLEASE NOTE

Begonia Tubers available January to March.
Begonia Seedlings available end May-June.
Begonia Seeds available December—
Montbretia Tubers available December to March.
Delphinium Plants available March-June and Sept.-November.
Delphinium Seed available fresh crop July.

Primrose Plants available August, September. Specimen Divisions.

Primrose Seedlings available February and March.

Primrose Seed available fresh crop July. Limited supply.

TUBERS and PLANTS are charged as a primarily of the United States.

TUBERS and PLANTS are charged at unit prices if labelled separately as to color etc. If ordered at dozen rate will be shipped assorted.

In the event of any enquiry concerning your order, kindly give fullest particulars possible, as with our hundreds of orders this will help us and expedite replies.

All stock guaranteed true to description but no guarantee for the result of growing over which we have no control.

CANADIAN CUSTOMERS will kindly procure permit from the Department of Agriculture, Ottawa, before Tubers are imported. No permit required for seed.

### **ANEMONES**

(St. Brigid—Anderson Strain)

We have the pleasure of distributing the above-named strain, which is a distinct improvement on the types offered generally in commerce—not only in wider range of color but also much larger flowers, carried on stiff, strong stems. In warm regions Anemones can be had in bloom from November to end of March by planting from July to October. In colder regions early spring planting is preferable.

PRICE OF COMPLETE MIXTURE . . . Large Tubers, \$1.00 per dozen; \$7.00 per 100. Medium size, 75c per dozen; \$5.00 per 100. (Orders of six tubers supplied at dozen rate; orders of fifty tubers supplied at 100 rate.)

**SEED**... Anemones can be grown easily from seed and will bloom four to five months after sowing. Sowing time in California, from May to October. In regions with cold winters, early spring only. **PRICE**... 50c per packet.

### **FREESIAS**

We had no intention of introducing our strain of Freesias until the named seedlings were sufficiently propagated for distribution. However, as we grow some 100,000 seedlings annually, we are in a position to offer mixtures of individual colors which will far outrank the standard named varieties of today. Each generation brings higher development and the seedling bulbs we are offering will produce very large flowers on tall, stiff stems which will reach, under glass, a height of from 20 to 30 inches. The color range covers many new shades not previously seen in Freesias. Their size and taller growth makes them a desirable subject for cut flower purposes during winter months. Planting time, September-October. From July to October we offer a complete mixture from our finest selection, covering the entire range of colors so far developed.

**PRICE OF COMPLETE MIXTURE** . . . Large bulbs, 75c per dozen; \$5.00 per 100. Medium size, 50c per dozen; \$3.00 per 100. (Orders of six tubers supplied at dozen rate; orders of fifty tubers supplied at 100 rate.)

**SEED** . . . Freesias can be grown from seed and bloom within six months of their sowing. The seed is comparatively large and germinates after thirty days. By sowing at intervals from May to August, one can have flowers from November to the end of March.

**PRICE** . . . 50c per packet of approximately 120 seeds; \$7.50 per ounce.



### **NERINE**

Planting time August-September . . . This charming South African member of the Amaryllis family is little known in this country. The brilliant color and beautiful form of its flowers which are produced in umbels during September and October, combined with simplicity of culture will endear it to all gardeners who cherish the rare and beautiful. We have imported all the new hybrids existing presently in Europe, but as the stock is very scarce we offer only one variety in quantity at present.

Nerine—Fothergilli Major . . . The king of all species. Brilliant orange scarlet with a gold-dust texture shimmering in sunlight.


**PRICE**... Strong bulbs from June to September. \$1.00 each; \$10.00 per dozen.

### **CULTURAL DIRECTIONS**

**SOIL** . . . After receiving the bulbs, pot up in six-inch pots in a mixture of one-third coarse leafmold, one-third loam with a little addition of sand and bonemeal. Plant the bulbs so that the upper part is exposed out of the ground, press the soil down firmly and water well. Do not water again until signs of growth show but keep fairly moist from then on.

**GROWING CONDITIONS** . . . Keep outdoors in full sunlight until danger of frost, then move either into the cool greenhouse or house where they will continue growing during the winter months. As soon as the weather permits, again take outdoors and by the end of May when the foliage gets yellow, dry off entirely.

**IMPORTANT** . . . Keep absolutely dry during the summer months as otherwise they would not produce flowers. By the first of September give a slight watering and when the buds begin to appear remove the upper part of the soil from the pot, taking care not to disturb the roots, and fill up with a new mixture of soil to which may be added either bonemeal or any complete commercial fertilizer. They will not need reporting and may be grown in the same pots for several years if they are given the dressing with rich soil each year. Their flowering period is September and October, after which the growth commences.



DEC 7 - 1940 \*